

Review Article

Impact of the essential oils on micro-organisms in meat preservation

Fahim A. Shaltout*Food Control Department, Faculty of Veterinary Medicine, Benha University, Egypt.***Corresponding Author:** Fahim A. Shaltout, Food Control Department, Faculty of Veterinary Medicine, Benha University, Egypt.**Received:** 📅 2024 Mar 23**Accepted:** 📅 2024 Apr 13**Published:** 📅 2024 Apr 22

Abstract

The Producers and the manufacturers have been challenged by the increasing demand for the safe and high-quality meat and meat products over the past few decades. Particularly, the recent demand for minimally the processed, the easily prepared, and the ready-to-eat meat products combined with the novel concepts of all-natural and clean-label has rapidly increased. These products may contain natural or organic ingredients without artificial preservatives that do not trigger the common food allergies or the sensitivities. The meat and the meat products are highly prone to microbial contamination since they are rich in essential nutrients and perishable. This is further accelerated by some intrinsic factors including pH and water activity of the fresh meat. In general, the freshest meat has a water activity value higher than 0.85, and its pH value falls within the favorable pH range for spoilage bacteria of the meat. Hence, deterioration in quality and potential public health issues is common if these products are not properly handled and preserved. The significant spoilage of the meat and the meat products occurs every year at different levels of the production chain including the preparation, the storage, and the distribution. Besides the lipid oxidation and the autolytic enzymatic spoilage, the microbial spoilage plays a significant role in this deterioration process leading to a substantial economic and environmental impact.

Keywords: Essential oils, safe, high quality, meat, meat products.

1. Introduction

The microorganisms associated with the spoilage of the meat and the meat products including bacteria such as the *Pseudomonas*, the *Acinetobacter*, the *Brochothrix thermosphacta*, the *Lactobacillus* spp., the *Enterobacter*, etc., and the yeast and the mold cause the quality defects such as of flavor, off-odor, etc. Additionally, the foodborne diseases have emerged as important and growing public health and economic problems in many countries over the last few decades [1-6]. The foodborne diseases are not limited to a particular age group or the country. The foodborne diseases occur each year resulting in the illnesses, with hospitalizations and deaths, leading to high medical costs and productivity losses. In regarding to the meat and the meat products, several pathogenic microorganisms including the *Salmonella* spp., the *Campylobacter jejuni*, the *Escherichia coli* O157:H7, the *Listeria monocytogenes*, the *Clostridium* spp. and the *Aeromonas hydrophila* can result in the foodborne illnesses to the consumers if the products are not preserved and handled properly [7-12]. The vegetative cells of the microbes are destroyed with the thermal processing, but not the spores of the food pathogens such as the *Bacillus* spp. and *Clostridium* spp. They can survive and need much harsher conditions to be inactivated. With the absence of the competitive microflora, these spores can germinate and grow under favorable

conditions caused by mishandling of the heat-treated products. Hence, it is vital to apply other hurdles such as low-temperature storage to avoid this problem [13-19]. To extend the period of the refrigerated storage, which is the most common method used for preserving the fresh meat and the meat products, many synthetic additives have been used over the years. The synthetic additives have been accused for some carcinogenic and toxic properties. This increased the consumer concerns towards the healthier meat and the meat products and the demand for the natural food additives over the years, which led researchers to examine the natural alternatives to the synthetic food additives [20-143]. The meat and the meat products are highly subject to the microbial deterioration, which ultimately leads to the safety and the quality issues if the meat are not properly handled and preserved. Several plant-derived Essential oils can be effectively used in the meat and the meat products as natural alternatives to the synthetic food additives, particularly as the effective antimicrobial agents [144-150]. The Phenolic compounds, such as the carvacrol, the eugenol, and the thymol, are mainly responsible for the antimicrobial activity of the Essential oils to increase the permeability of cell membranes and leading to loss of the cellular constituents. These natural additives should improve the meat quality without leaving residues in the product or the environment [20-26].

1.1. The Microbial deterioration of the meat and the meat products

Depending on the preservation method used, the growth and the metabolism of the spoilage and the pathogenic microorganisms can cause rapid spoilage of the meat and the meat products and the serious foodborne intoxications. The Bacteria are generally considered as the principle agents responsible for these deteriorations and the health issues in addition to the molds and yeasts presents the major genera of bacteria, yeasts, and molds found in meat and meat products [27-33]. Main spoilage bacteria including the *Pseudomonas*, the *Acinetobacter*, the *B. thermosphacta*, the *Moraxella*, the *Enterobacter*, the *Lactobacillus* spp., the *Leuconostoc* spp., the *Proteus* spp. etc, the yeast and the mold, the decompose meat and the meat products and develop the unpleasant quality characteristics when they grow in large numbers in these perishable products [34-40]. This can be attributed to the degradation of the proteins and the lipids present in the meat and the meat products resulting in the off-odors, the off-flavors, the discoloration, the texture defects, the slime and the gas productions, and the changes in pH. Although the spoilage microorganisms normally do not cause the illness, they can result in the gastrointestinal disturbances when consumed in the high concentrations [41-48]. However, the rate of the meat spoilage is affected by several factors including the hygiene, the storage temperature, and the acidity of the meat and the meat products. The growth of the spoilage bacteria is favored at a pH range of 5.5-7.0. The *Campylobacter jejuni*, *Salmonella* serotype Typhimurium, *Escherichia coli* O157:H7, other Enterohemorrhagic *E. coli* (EHEC), *Listeria monocytogenes*, *Arcobacter butzleri*, *Mycobacterium avium* subsp. *paratuberculosis* and *Aeromonas hydrophila* are the most prevalent and serious emerging pathogens in the meat and the meat-derived products [49-56]. The color, the odor, the taste or the texture of the meat and the meat products are not often changed by these pathogenic bacteria. which are mainly responsible for the food poisoning and the food intoxications. The microbial deterioration of the meat and the meat products can be considered as one of the main limitations in the meat industry. The proper protective practices should be, therefore, applied for the meat and the meat products to produce the highest quality products possible [57-64]. The common preservation methods used against the microbial deterioration Due to the increasing demand for the precooked, the refrigerated, the ready-to-eat meat, and the meat products convenient for the modern and the busy lifestyles, the meat manufacturers must overcome many challenges, including the control of the microbial deterioration of these products [65-72]. Several factors such as the microbial growth, the color, and the lipid oxidation affect the shelf life and the consumer acceptance of the fresh meat and the meat products. Therefore, to produce the safest and the highest quality products possible, these factors must be controlled [73-79]. Several of the thermal and the non-thermal food preservation techniques have been used, alone or in combinations, to prevent or minimize the growth of spoilage and the pathogenic microorganisms in the meat and the meat products. The common physical and the chemical preservation techniques used to control the bacterial activity [80-87].

1.2. The mode of action of the Essential oils

The antimicrobial activity of a given Essential oil can be attributed to its major constituents as well as their interaction with the minor constituents present in oils. However, the antimicrobial activity of the Essential oils has been consistently linked to phenolic constituents such as the carvacrol, the eugenol, and the thymol. The presence of the hydroxyl groups in the phenolic compounds is very vital for their antimicrobial activity [88-94]. The antimicrobial activity of the Essential oils is not attributable to one specific mechanism. Several locations or mechanisms in the microbial cells are supposed to be the sites of action for the Essential oil constituents. In brief, Essential oils can degrade the cell wall, disturb the phospholipid bilayer of the cytoplasmic membrane, and damage the membrane proteins leading to increased permeability of the cell membrane and loss of cellular constituents [96-102]. The Essential oils can further disrupt the proton motive force, electron flow, and active transport, and coagulate the cell contents. The Essential oils can impair a variety of enzyme systems including the enzymes involved in the energy regulation and synthesis of structural components and inactivate or destroy genetic material, strengthening their antimicrobial activities [103-109].

1.3. Limitations and Future Perspectives

Although good antimicrobial activities were observed for many Essential oils, some limitations have also been identified in the application of the Essential oils in the meat and the meat products. The interaction of some Essential oils with the food ingredients and structure may decrease their effectiveness [110-116]. The markedly reduced activity of the Essential oils may result in the food systems such as the meat and the meat products when compared to in vitro results. This may be attributed to the presence of fats, carbohydrates, proteins, and salts in such systems. For instance, the mint and the cilantro Essential oils were not effective against the *L. monocytogenes* in the products containing high levels of the fat, such as the pâté and a coating for the ham containing the canola oil [117-121]. It may be difficult to maintain the quality consistency because the composition of an individual Essential oil can vary due to several factors including the time of harvesting, variety, the part of the plant used, and method of extraction. The antimicrobial potency of the Essential oil constituents depends on pH, temperature, and level of microbial contamination [123-129]. Further, use of the Essential oils as the preservatives in the food has been limited as they are required in high concentrations in order to achieve the sufficient antimicrobial activity. The use of some Essential oils is mainly restricted to the spicy foods including the meat products that are associated with the herbs, spices, or seasonings since they affect the organoleptic quality of the food [130-136].

2. Conclusion

The application of the Essential oils is partially limited due to their intense aroma which may cause negative organoleptic effects. Novel technologies such as encapsulation of the Essential oils into the Nano emulsions and the use of the Essential oils as part of the hurdle technology (combined processes with MAP, nisin, EDTA, lysozyme etc.) to improve

the microbial stability and the sensory quality of the meat and the meat products are being used in the meat industry; traditional methods of adding the Essential oils directly into the meat batter during manufacturing of the meat products.

Conflicts of Interest

The author declares no conflicts of interest regarding the publication of this paper.

References

- Nataro, J. P., & Kaper, J. B. (1998). Diarrheagenic *Escherichia coli*. *Clinical microbiology reviews*, 11(1),142-201.
- Shaltout, F. A., Riad, E. M., & Asmaa, A. E. PREVALENCE OF MYCOBACTERIUM SPP. IN CATTLE MEAT AND OFFAL'S SLAUGHTERED IN AND OUT ABATTOIR.
- McDonnell, M. J., Rivas, L., Burgess, C. M., Fanning, S., & Duffy, G. (2012). Evaluation of carvacrol for the control of *Escherichia coli* O157 on cattle hide and carcass cuts. *Foodborne Pathogens and Disease*, 9(11),1049-1052.
- Abd Elaziz, O., Hassanin, F., Shaltout, F., & Mohammed, O. (2021). Prevalence of some foodborne parasitic affection in slaughtered animals in local Egyptian abattoir. *Benha Veterinary Medical Journal*, 41(1), 111-114.
- Cornforth, D. (1994). Color—its basis and importance. In *Quality attributes and their measurement in meat, poultry and fish products* (pp. 34-78). Boston, MA: Springer US.
- El-Aziz, A., Mohamed, O., Hassanin, F. S., Shaltout, F. A. E., & Mohammed, O. (2021). Prevalence of some zoonotic parasitic affections in sheep carcasses in a local abattoir in Cairo, Egypt. *Benha Veterinary Medical Journal*, 41(1), 115-119.
- Fasseas, M. K., Mountzouris, K. C., Tarantilis, P. A., Polissiou, M., & Zervas, G. (2008). Antioxidant activity in meat treated with oregano and sage essential oils. *Food chemistry*, 106(3),1188-1194.
- Al Shorman, A.A.M. ;Shaltout, F.A. and hilat, N (1999): Detection of certain hormone residues in meat marketed in Jordan. *Jordan University of Science and Technology*, 1st International Conference on Sheep and goat Diseases and Productivity, 23-25 October, 1999.
- Souza, A. A., Dias, N. A. A., Piccoli, R. H., & Bertolucci, S. K. V. (2016). Composição química e concentração mínima bactericida de dezesseis óleos essenciais sobre *Escherichia coli* enterotoxigênica. *Revista Brasileira de Plantas Mediciniais*, 18, 105-112.
- Saleh, E., Shaltout, F., & Abd Elaal, E. (2021). Effect of some organic acids on microbial quality of dressed cattle carcasses in Damietta abattoirs, Egypt. *Damanhour Journal of Veterinary Sciences*, 5(2), 17-20.
- Tserennadmid, R., Takó, M., Galgóczy, L., Papp, T., Vágvolgyi, C., Gerő, L., & Krisch, J. (2010). Antibacterial effect of essential oils and interaction with food components. *Open Life Sciences*, 5(5), 641-648.
- Shaltout, F. A. E. (2017). Microbiological evaluation of some heat treated fish products in Egyptian markets. *EC Nutrition*, 12, 124-132.
- Chorianopoulos, N., Kalpoutzakis, E., Aligiannis, N., Mitaku, S., Nychas, G. J., & Haroutounian, S. A. (2004). Essential oils of *Satureja*, *Origanum*, and *Thymus* species: chemical composition and antibacterial activities against foodborne pathogens. *Journal of Agricultural and Food Chemistry*, 52(26),8261-8267.
-
- Edris, A., Hassan, M., Saltout, F., & El-Hosseney, S. (2013). Chemical evaluation of cattle and camel meat. *BVMJ*, 25, 145-150.
- Hayouni, E. A., Chraief, I., Abedrabba, M., Bouix, M., Leveau, J. Y., Mohammed, H., & Hamdi, M. (2008). Tunisian *Salvia officinalis* L. and *Schinus molle* L. essential oils: Their chemical compositions and their preservative effects against *Salmonella* inoculated in minced beef meat. *International Journal of Food Microbiology*, 125(3), 242-251.
- Edris, A., Hassan, M., Shaltout, F., & El-Hosseney, S. (2012). Detection of *E. coli* and *Salmonella* organisms in cattle and camel meat. *Benha veterinary medical journal*, 24(2), 198-204.
- Govaris, A., Solomakos, N., Pexara, A., & Chatzopoulou, P. S. (2010). The antimicrobial effect of oregano essential oil, nisin and their combination against *Salmonella* Enteritidis in minced sheep meat during refrigerated storage. *International journal of food microbiology*, 137(2-3), 175-180.
- Edris, A. M., Hemmat, M. I., Shaltout, F. A., Elshater, M. A., & Eman, F. M. I. (2012). Study on incipient spoilage of chilled chicken cuts-up. *Benha Veterinar y*, 23, 81-86
- Ramos, E. M., & de Miranda Gomide, L. A. (2007). Avaliação da qualidade de carnes: fundamento e metodologias. *UFV*.
- Edris, A. M., Hemmat, M. I., Shaltout, F. A., Elshater, M. A., & Eman, F. M. I. (2012). Chemical analysis of chicken meat with relation to its quality. *Benha veterinary medical Journal*, 23(1), 87-93.
- Van Den Dool, H. A. N. D., & Kratz, P. D. (1963). A generalization of the retention index system including linear temperature programmed gas-liquid partition chromatography. *Journal of chromatography*.
- Edris, A. M., Shaltout, F. A., & Abd Allah, A. M. (2005). Incidence of *Bacillus cereus* in some meat products and the effect of cooking on its survival. *Zag. Vet. J*, 33(2), 118-124.
- Adams, R. P. Identification of Essential Oil Components by Gas Chromatography/Mass Spectroscopy.
- Edris, A. M., Shaltout, F. A., & Arab, W. S. (2005). Bacterial evaluation of quail Meat. *Benha Vet. Med. J*, 16(1), 1-14.
- Aziz, M., & Karboune, S. (2018). Natural antimicrobial/antioxidant agents in meat and poultry products as well as fruits and vegetables: A review. *Critical reviews in food science and nutrition*, 58(3), 486-511.
- Edris, A.M.; Shaltout, F.A. ;Salem, G.H. and El-Toukhy, E.I. (2011): Incidence and isolation of *Salmonellae* from some meat products. *Benha University ,Faculty of Veterinary Medicine , Fourth Scientific Conference 25-27th May 2011 Veterinary Medicine and Food Safety)* 172-179 benha , Egypt.
- Souza, A. A. (2015). Análise química e potencial anti-

- crobiano de óleos essenciais sobre *Escherichia coli* enterotóxigênica na conservação de carne moída.
29. Shaltout, F. A. E. (2017). Microbiological evaluation of some heat treated fish products in Egyptian markets. *EC Nutrition*, 12, 124-132.
 30. Banwart, G. J. (1989). *Basic food microbiology*.
 31. Edris, A. M., Shaltout, F. A., Salem, G. H., & El-Toukhy, E. I. (2011). Incidence and isolation of *Salmonellae* from some meat products. Benha University, Faculty of Veterinary Medicine. In Fourth Scientific Conference (pp. 172-179).
 32. Burt, S. (2004). Essential oils: their antibacterial properties and potential applications in foods—a review. *International journal of food microbiology*, 94(3), 223-253.
 33. Ragab, A., Edris, A. M., Shaltout, F. A., & Salem, A. M. (2022). Effect of titanium dioxide nanoparticles and thyme essential oil on the quality of the chicken fillet. *Benha Veterinary Medical Journal*, 41(2), 38-40.
 34. de Oliveira, T. L. C., de Araújo Soares, R., & Piccoli, R. H. (2013). A Weibull model to describe antimicrobial kinetics of oregano and lemongrass essential oils against *Salmonella* Enteritidis in ground beef during refrigerated storage. *Meat science*, 93(3), 645-651.
 35. Hassan, M. A., Shaltout, F. A., Arfa, M. M., Mansour, A. H., & Saudi, K. R. (2013). Biochemical studies on rabbit meat related to some diseases. *Benha Vet. Med. J*, 25(1), 88-93.
 36. Gill, A. O., Delaquis, P., Russo, P., & Holley, R. A. (2002). Evaluation of antilisterial action of cilantro oil on vacuum packed ham. *International journal of food microbiology*, 73(1), 83-92.
 37. MA, H., & FA, S. (1997). Occurrence of some food poisoning microorganisms in rabbit carcasses.
 38. Cho, S. H., Kim, J., Oh, K. H., Hu, J. K., Seo, J., Oh, S. S., ... & Choe, Y. J. (2014). Outbreak of enterotoxigenic *Escherichia coli* O169 enteritis in schoolchildren associated with consumption of kimchi, Republic of Korea, 2012. *Epidemiology & Infection*, 142(3), 616-623.
 39. Hassan, M., Shaltout, F. A., & Saqur, N. (2020). Histamine in some fish products. *Archives of Animal Husbandry & Dairy Science*, 2(1), 1-3.
 40. von Mentzer, A., Connor, T. R., Wieler, L. H., Semmler, T., Iguchi, A., Thomson, N. R., ... & Dougan, G. (2014). Identification of enterotoxigenic *Escherichia coli* (ETEC) clades with long-term global distribution. *Nature genetics*, 46(12), 1321-1326.
 41. Hassan, M. A., Shaltout, F. A., Arfa, M. M., Mansour, A. H., & Saudi, K. R. (2013). Biochemical studies on rabbit meat related to some diseases. *Benha Vet. Med. J*, 25(1), 88-93.
 42. Jayari, A., El Abed, N., Jouini, A., Mohammed Saed Abdul-Wahab, O., Maaroufi, A., & Ben Hadj Ahmed, S. (2018). Antibacterial activity of *Thymus capitatus* and *Thymus algeriensis* essential oils against four food-borne pathogens inoculated in minced beef meat. *Journal of Food Safety*, 38(1), e12409.
 43. Hassan, M. A., Shaltout, F. A., Maarouf, A. A., & El-Shafey, W. S. (2014). Psychrotrophic bacteria in frozen fish with special reference to *Pseudomonas* species. *Benha Vet. Med. J*, 27(1), 783.
 44. Pateiro, M., Barba, F. J., Domínguez, R., Sant'Ana, A. S., Khaneghah, A. M., Gavahian, M., ... & Lorenzo, J. M. (2018). Essential oils as natural additives to prevent oxidation reactions in meat and meat products: A review. *Food research international*, 113, 156-166.
 45. Hassan, M. A., Shaltout, F. A., Arafa, M. M., Mansour, A. H., & Saudi, K. R. (2013). Bacteriological studies on rabbit meat related to some diseases. *Benha Vet Med J*, 25(1), 94-99.
 46. Smith-Palmer, A., Stewart, J., & Fyfe, L. (2001). The potential application of plant essential oils as natural food preservatives in soft cheese. *Food microbiology*, 18(4), 463-470.
 47. Hassanin, F. S., Hassan, M. A., Shaltout, F. A., Shawqy, N. A., & Abd-Elhameed, G. A. (2017). Chemical criteria of chicken meat. *Benha Veterinary Medical Journal*, 33(2), 457-464.
 48. Dal Pozzo, M., Viégas, J., Santurio, D. F., Rossatto, L., Soares, I. H., Alves, S. H., & Costa, M. M. D. (2011). Atividade antimicrobiana de óleos essenciais de condimentos frente a *Staphylococcus* spp isolados de mastite caprina. *Ciência Rural*, 41, 667-672.
 49. Hassanien-faten, S., Hassan, M. A., Shaltout, S., & El-raï-Amina, M. CLOSTRIDIUM PERFRINGENS IN VACUUM PACKAGED MEAT PRODUCTS.
 50. Jayasena, D. D., & Jo, C. (2013). Essential oils as potential antimicrobial agents in meat and meat products: A review. *Trends in Food Science & Technology*, 34(2), 96-108.
 51. Hassanien, F., Shaltout, F. A. E., Fahmey, M. Z., & Elsukkary, H. F. A. (2020). Bacteriological quality guides in local and imported beef and their relation to public health. *Benha Veterinary Medical Journal*, 39(1), 125-129.
 52. Hassanien, F., Shaltout, F. A. E., Fahmey, M. Z., & Elsukkary, H. F. A. (2020). Bacteriological quality guides in local and imported beef and their relation to public health. *Benha Veterinary Medical Journal*, 39(1), 125-129.
 53. Hassanin, F. S., Shaltout, F. A., & Afifi, M. E. (2013). Parasitic affections in edible offal. *Benha Vet. Med. J*, 25(1), 46-55.
 54. Bakkali, F., Averbeck, S., Averbeck, D., & Idaomar, M. (2008). Biological effects of essential oils—a review. *Food and chemical toxicology*, 46(2), 446-475.
 55. Hassanin, F. S., Shaltout, F. A., Lamada, H. M., & Abd Allah, E. M. (2011). The effect of preservative (nisin) on the survival of *Listeria monocytogenes*. *Benha Veterinary Medical Journal*, 141-145.
 56. Nazzaro, F., Fratianni, F., De Martino, L., Coppola, R., & De Feo, V. (2013). Effect of essential oils on pathogenic bacteria. *Pharmaceuticals*, 6(12), 1451-1474.
 57. Khattab, E., Shaltout, F., & Sabik, I. *Benha Veterinary Medical Journal*.
 58. Höferl, M., Buchbauer, G., Jirovetz, L., Schmidt, E., Stoyanova, A., Denkova, Z., ... & Geissler, M. (2009). Correlation of antimicrobial activities of various essential oils and their main aromatic volatile constituents. *Journal of Essential Oil Research*, 21(5), 459-463.
 - 59.
 60. Saad, S. M., Shaltout, F. A., Farag, A. A., & Mohammed, H. F. (2022). Organophosphorus Residues in Fish in Rural Areas. *Journal of Progress in Engineering and Physical Science*, 1(1), 27-31.

61. Saif, M., Saad, S., Shaltout, F., Hassanin, F. S., & Zaghloul, M. (2019). Molecular detection of enterotoxigenic *Staphylococcus aureus* in ready to eat beef products. *Benha Veterinary Medical Journal*, 37(1), 7-11.
62. Aureli, P., Costantini, A., & Zolea, S. (1992). Antimicrobial activity of some plant essential oils against *Listeria monocytogenes*. *Journal of food protection*, 55(5), 344-348.
63. Saif, M., Saad, S., Hassanin, F. S., Shaltout, F., & Zaghloul, M. (2019). Prevalence of methicillin-resistant *Staphylococcus aureus* in some ready-to-eat meat products. *Benha Veterinary Medical Journal*, 37(1), 12-15.
64. Badi, H. N., Yazdani, D., Ali, S. M., & Nazari, F. (2004). Effects of spacing and harvesting time on herbage yield and quality/quantity of oil in thyme, *Thymus vulgaris* L. *Industrial crops and products*, 19(3), 231-236.
65. Abdelmordy, A., Shaltout, F., & Saad, S. M. (2023). Studies on Pesticides Residues in Fish in Menofia Governorate. *Benha Journal of Applied Sciences*, 8(5), 323-330.
66. Al-Reza, S. M., Rahman, A., Lee, J., & Kang, S. C. (2010). Potential roles of essential oil and organic extracts of *Zizyphus jujuba* in inhibiting food-borne pathogens. *Food Chemistry*, 119(3), 981-986.
67. Abdelmordy, A., Shaltout, F., & Saad, S. M. (2023). Organochlorine Residues in Fish in Rural Areas. *Benha Journal of Applied Sciences*, 8(5), 331-336.
68. Ayala-Zavala, J. F., Oms-Oliu, G., Odriozola-Serrano, I., González-Aguilar, G. A., Álvarez-Parrilla, E., & Martín-Belloso, O. (2008). Bio-preservation of fresh-cut tomatoes using natural antimicrobials. *European Food Research and Technology*, 226, 1047-1055.
69. Shaltout, F., Hussein, M., & khaled Elsayed, N. (2023). Histological Detection of Unauthorized Herbal and Animal Contents in Some Meat Products. *Journal of Advanced Veterinary Research*, 13(2), 157-160.
70. Ghanem, A., Shaltout, F., & Heikal, G. I. (2022). Mycological quality of some chicken meat cuts in Gharbiya governorate with special reference to *Aspergillus flavus* virulent factors. *Benha Veterinary Medical Journal*, 42(1), 12-16.
71. Darwish, W., Shaltout, F. A., Salem, R. M., Eldiasty, E. M., & Diab, F. A. (2022). Seasonal Impact on the Prevalence of Yeast Contamination of Chicken Meat Products and Edible Giblets. *Journal of Advanced Veterinary Research*, 12(5), 641-644.
72. Shaltout, F. A., Barr, A. A. H., & Abdelaziz, M. E. (2022). Pathogenic Microorganisms in Meat Products. *Biomedical Journal of Scientific & Technical Research*, 41(4), 32836-32843.
73. Shaltout, F. A., & Koura, H. A. (2017). Impact of some essential oils on the quality aspect and shelf life of meat. *Benha Veterinary Medical Journal*, 33(2), 351-364.
74. Mohammed, I. Z. (2020). Bacteriological profile of some raw chicken meat cuts in Ismailia city, Egypt. *Benha Veterinary Medical Journal*, 39(1), 11-15.
75. Afify, S. A., Shaltout, F., & Mohammed, I. Z. (2020). Detection of *E. coli* O157 and *Salmonella* species in some raw chicken meat cuts in Ismailia province, Egypt. *Benha Veterinary Medical Journal*, 39(1), 101-104.
76. Shaltout, F. A., El-diasty, E. M., & Asmaa-Hassan, M. A. (2019). HYGIENIC QUALITY OF READY TO EAT COOKED MEAT IN RESTAURANTS AT CAIRO GOVERNORATE. *Journal of Global Biosciences*, 8(12), 6627-6641.
77. Shaltout, F. A., Nasief, M. Z., Lotfy, L. M., & Gamil, B. T. *Benha Veterinary Medical Journal*.
78. Shaltout, F.A.(2019): Poultry Meat. *Scholarly Journal of Food and Nutrition* 22 1-2.
79. Saad, S. M., Shaltout, F. A., Abou Elroos, N. A., & El-nahas, S. B. (2019). Antimicrobial effect of some essential oils on some pathogenic bacteria in minced meat. *Journal of Food Science and Nutrition Research*, 2(1), 13-21.
80. Arakeeb, S. M., Hassanien, F., Shaltout, F., & Homouda, S. (2019). Natural preservatives in raw chicken meat. *Benha Veterinary Medical Journal*, 37(1), 41-45.
81. Hazaa, W., Shaltout, F., & El-Shater, M. A. (2019). Prevalence of some chemical hazards in some meat products. *Benha Veterinary Medical Journal*, 37(1), 32-36.
82. Hazaa, W., Shaltout, F., & El-Shater, M. (2019). Identification of some biological hazards in some meat products. *Benha Veterinary Medical Journal*, 37(1), 27-31.
83. Gaafar, R., Hassanin, F. S., Shaltout, F., & Zaghloul, M. (2019). Molecular detection of enterotoxigenic *Staphylococcus aureus* in some ready to eat meat based sandwiches. *Benha Veterinary Medical Journal*, 37(1), 22-26.
84. Gaafar, R., Hassanin, F. S., Shaltout, F., & Zaghloul, M. (2019). Hygienic profile of some ready to eat meat product sandwiches sold in Benha city, Qalubiyah Governorate, Egypt. *Benha Veterinary Medical Journal*, 37(1), 16-21.
85. Saad, S. M., Shaltout, F. A., Abou Elroos, N. A., & El-nahas, S. B. (2019). Antimicrobial effect of some essential oils on some pathogenic bacteria in minced meat. *Journal of Food Science and Nutrition Research*, 2(1), 13-21.
86. Shaltout, F. A., Saad, M., Abo El-Roos, N., & Saber, E. (2019). Incidence of *Staphylococci* and *E. coli* in Meat and Some Meat Products. *EC Nutrition*, 14.
87. Shaltout, F. (2019). Pollution of Chicken Meat and Its Products by Heavy Metals. *Research and Reviews on Healthcare: Open Access Journal*, 4(3), 381-3382.
88. Shaltout, F. A., EL-diasty, E. M., & Mohamed, M. S. (2018). Effects of chitosan on quality attributes fresh meat slices stored at 40C. *Benha Veterinary Medical Journal*, 35(2), 157-168.
89. Shaltout, F. A., & Abdel-Aziz, A. M. (2004). *Salmonella enterica* serovar enteritidis in poultry meat and their epidemiology.
90. Shaltout, F. A., El Zahaby, D. I., Lotfy, L. M., & El-Shorah, H. F. (2018). Bacteriological profile of chicken meat products. *Food Nutr Current Res*, 1(3), 83-90.
91. Shaltout, F., El-Shater, M. A., El-Aziz, A., & Mohamed, W. (2015). Bacteriological assessment of street vend- ed meat products sandwiches in Kalyobia Governorate. *Benha Veterinary Medical Journal*, 28(2), 58-66.
92. Fahim, H. M. (2019). Evaluate antibiotic residues in beef and effect of cooking and freezing on it. *Benha Veterinary Medical Journal*, 36(2), 109-116.
93. Shaltout, F. A., Zakaria, I. M., & Nabil, M. E. (2018). Incidence of some anaerobic bacteria isolated from chicken meat products with special reference to *Clostridium perfringens*. *Nutrition and Food Toxicology*, 2(5), 429-438.
94. Shaltout, F. A., Maarouf, A. A., & Elkhouly, M. E. (2017).

- Bacteriological Evaluation of Frozen Sausage. *Nutrition and Food Toxicology*, 1(2017), 174-185.
95. Shaltout, F. A., El-Toukhy, E. I., & Abd El-Hai, M. M. (2019). Molecular Diagnosis of Salmonellae in Frozen Meat and Some Meat Products. *Nutrition Food Technol Open Access*, 5(1).
 96. Shaltout, F. A., Ali, A. M., & Rashad, S. M. (2016). Bacterial contamination of fast foods. *Benha Journal of Applied Sciences (BJAS)*, 1(2), 45-51.
 97. Shaltout, F. A., Eltanani, J., & Elmelegy, A. S. (2015). Microbiological status of meat and chicken received to University student hostel. *Benha Veterinary Medical Journal*, 29(2), 187-192.
 98. Saad, M. S., Edris, A. M., Shaltout, F. A., & Shima, E. (2011). Isolation and identification of Salmonellae and E. coli from meat and poultry cuts by using multiplex PCR. *Benha Veterinary Medical Journal*, 22(2), 152-160.
 99. Saad, S. M., & Shaltout, F. A. (1998). Mycological evaluation of camel carcasses at Kalyobia abattoirs.
 100. Saad, S. M., Shaltout, F. A., Abou Elroos, N. A., & El-nahas, S. B. (2019). Antimicrobial effect of some essential oils on some pathogenic bacteria in minced meat. *Journal of Food Science and Nutrition Research*, 2(1), 13-21.
 101. Saad S.M. , Hassanin, F. S; Shaltout, F.A., Marionette Z Nassif, Marwa Z Seif.(2019): Prevalence of Methicillin-Resistant Staphylococcus Aureus in Some Ready-to-Eat Meat Products. *American Journal of Biomedical Science & Research* 4(6):460-464.
 102. Shaltout, F. A., Saad, M., Abo El-Roos, N., & Saber, E. (2019). Incidence of Staphylococci and E. coli in Meat and Some Meat Products. *EC Nutrition*, 14.
 103. Ahmed, T. (2017). Studying the effect of gamma irradiation on bovine offal's infected with Mycobacterium tuberculosis bovine type. *Veterinary Medical Journal (Giza)*, 63(4), 17-24.
 104. Shaltout, F. A., Zakaria, I. M., & Nabil, M. E. (2018). Incidence of some anaerobic bacteria isolated from chicken meat products with special reference to Clostridium perfringens. *Nutrition and Food Toxicology*, 2(5), 429-438.
 105. Faheim, A. S., Mohamed, A. H., & Fatin, S. H. THERMAL INACTIVATION OF ENTEROHAEMORRHAGIC ESCHERICHIA COLI O157: H7 AND ITS SENSITIVITY TO NISIN AND LACTIC ACID CULTURES.
 106. Shaltout, F. A., El-diasty, E. M., Elmesalmy, M., & Elshaer, M. (2014). Study on fungal contamination of some chicken meat products with special reference to 2 the use of PCR for its identification. In *Conf. Vet. Med. J. Giza (Vol. 60, pp. 1-10)*.
 107. Shaltout, F. A. (2002). Microbiological aspects of semi-cooked chicken meat products. *Benha Vet. Med. J*, 13(2), 15-26.
 108. Shaltout, F. A., & Koura, H. A. (2017). Impact of some essential oils on the quality aspect and shelf life of meat. *Benha Veterinary Medical Journal*, 33(2), 351-364.
 109. Shaltout, F. A. E., Farouk, M., Ibrahim, H. A., & Afifi, M. E. (2017). Incidence of Coliform and Staphylococcus aureus in ready to eat fast foods. *Benha Veterinary Medical Journal*, 32(1), 13-17.
 110. Shaltout, F. A., Zakaria, I. M., & Nabil, M. E. (2017). Detection and typing of Clostridium perfringens in some retail chicken meat products. *BMVJ*, 33, 283-291.
 111. Shaltout, F. A. M. (1992). Studies on mycotoxins in meat and meat by-products.
 112. Shaltout, F. A. M. (1996). Mycological and mycotoxicological profile of some meat products.
 113. Shaltout, F. A. (1998). Proteolytic Psychrotrophes in Some Meat products. *Alex. Vet. Med. J*, 14(2), 97-107.
 114. Shaltout, F. A. (1999). Anaerobic bacteria in vacuum packed meat products. *Benha Vet. Med. J*, 10(1), 1-10.
 115. Shaltout, F. A. (2000). PROTOZOAL F OODBORNE PATHOGENS IN SOME MEAT PRODCUTS. *Assiut Veterinary Medical Journal*, 42(84), 54-59.
 116. Shaltout, F. A. (2001). Quality evaluation of sheep carcasses slaughtered at Kalyobia abattoirs. *Assiut Veterinary Medical Journal*, 46(91), 150-159.
 117. Shaltout, F. A. (2002). Microbiological aspects of semi-cooked chicken meat products. *Benha Vet. Med. J*, 13(2), 15-26.
 118. Shaltout, F. A. (2003, April). Yersinia Enterocolitica in some meat products and fish marketed at Benha city. In *The Third international conference Mansoura (pp. 29-30)*.
 119. Fahim, A. S. (2020). Microbiological Quality of Chicken Carcasses at Modern Poultry Plant. *J. Nutrition and Food Processing*, 3(1).
 120. Shaltout, F. A., & Abdel-Aziz, A. M. (2004). Salmonella enterica serovar enteritidis in poultry meat and their epidemiology.
 121. Shaltout, F. A., & Abdel Aziz, A. M. (2004). ESCHERICHIA COLI STRAINS IN SLAUGHTERED ANIMALS AND THEIR PUBLIC HEALTH IMPORTENCE. *J. Egypt. Vet. Med. Association*, 64(2), 7-21.
 122. Shaltout, F. A., Amin, R. A., Nassif, M. Z., & Abd-Elwahab, S. A. (2014). Detection of aflatoxins in some meat products. *Benha Vet Med J*, 27(2), 368-74.
 123. Shaltout, F. A., Afify, J. R. E., & Abo Elhasan, A. A. (2012). Improvement of microbiological status of oriental sausage. *Journal of Egyptian Veterinary Medical Association*, 72(2), 157-167.
 124. Shaltout, F. A. (2023). Chemical analytical studies on rabbit meat and liver. *Benha Veterinary Medical Journal (Egypt)*, 7(2).
 125. Shaltout, F. A., & Edris, A. M. (1999). Contamination of shawerma with pathogenic yeasts. *Assiut Veterinary Medical Journal*, 41(81), 170-176.
 126. Shaltout, F. A., El-diasty, E. M., & Mohamed, M. S. (2014, June). Incidence of lipolytic and proteolytic fungi in some chicken meat products and their public health significance. In *Animal Health Research Institute: First International Conference on Food Safety and Technology (pp. 19-23)*.
 127. Shaltout, F. A., El-diasty, E. M., Salem, R. M., & Asmaa, M. A. (2016). Mycological quality of chicken carcasses and extending shelf-life by using preservatives at refrigerated storage. *Veterinary Medical Journal-Giza*, 62(3), 1-10.
 128. Shaltout, F. A., Salem, R., Eldiasty, E. M., & Diab, F. (2016). Mycological evaluation of some ready to eat meat products with special reference to molecular char-

- acterization. *Vet. Med. J. Giza*, 62, 9-14.
129. Shaltout, F., El-Shater, M. A., El-Aziz, A., & Mohamed, W. (2015). Bacteriological assessment of street vended meat products sandwiches in Kalyobia Governorate. *Benha Veterinary Medical Journal*, 28(2), 58-66.
 130. Shaltout, F. A., Gerges, M. T., & Shewail, A. A. (2014). Impact of organic acids and their salts on microbial quality and shelf life of beef meat.
 131. Shaltout, F. A., Ghoneim, A. M., Essmail, M. E., & Yousseif, A. (2001). Studies on aflatoxin B1 residues in rabbits and their pathological effects. *J. Egypt. Vet. Med. Ass*, 61, 85-103.
 132. Shaltout, F. A., & Hanan, M. T. El-Laewndy,(2003): Heavy metal residues in Shawerma. *Beni-Sueif Vet. Med. J*, 12(1), 213-224.
 133. Shaltout, F. A. (2002). Microbiological aspects of semi-cooked chicken meat products. *Benha Vet. Med. J*, 13(2), 15-26.
 134. Fahim, S. A., Mohammed, H. F., & Saber, E. N. (2015). Detection of some heavy metals in fish (*Tilapia nilotica* and *Clarias lazera*) at Menofia governorate. *Benha Veterinary Medical Journal*, 29(1), 124-128.
 135. Shaltout, F. A., & Ibrahim, H. M. (1997). Quality evaluation of luncheon and Alexandrian sausage. *Benha Vet. Med. J*, 10(1), 1-10.
 136. Shaltout, F. A., Marionette, Z. N., & Shakran, A. M. (2014). Quality of battered and breaded chicken meat products.
 137. Shaltout, F. A., Amani, M. S., Mahmoud, A. H., & Abd Elraheem, K. A. (2013). Bacterial aspect of cooked meat and edible offal at street vendors level. *J. Benha vet. Med*, 24(1), 320-328.
 138. Shaltout, F. A., & Salem, R. M. T. (2000). Moulds, aflatoxin B1 and ochratoxin A in frozen livers and meat products.
 139. Al-Tarazi, Y., Al-Zamil, A., Shaltout, F., & Abdel-Samei, H. (2003). Sanitary status of raw cow milk marketed in northern Jordan. *Assiut Veterinary Medical Journal*, 49(96), 180-194.
 140. Shaltout, F. A., Zakaria, I. M., & Nabil, M. E. (2018). Incidence of some anaerobic bacteria isolated from chicken meat products with special reference to *Clostridium perfringens*. *Nutrition and Food Toxicology*, 2(5), 429-438.
 141. Shaltout, F. A., El-diasty, E. M., & Mohamed, M. S. (2014, June). Incidence of lipolytic and proteolytic fungi in some chicken meat products and their public health significance. In *Animal Health Research Institute: First International Conference on Food Safety and Technology* (pp. 19-23).
 142. Shaltout, F. A., El-diasty, E. M., Salem, R. M., & Asmaa, M. A. (2016). Mycological quality of chicken carcasses and extending shelf-life by using preservatives at refrigerated storage. *Veterinary Medical Journal-Giza*, 62(3), 1-10.
 143. Shaltout, F. A., Salem, R. M., El-Diasty, E. M., & Hassan, W. I. M. (2019). Effect of Lemon fruits and turmeric extracts on fungal pathogens in refrigerated chicken fillet meat. *Global Veterinaria*, 21(3), 156-160.
 144. Shaltout, F. A., El-diasty, E. M., Elmesalamy, M., & Elshaer, M. (2014). Study on fungal contamination of some chicken meat products with special reference to the use of PCR for its identification. In *Conf. Vet. Med. J. Giza* (Vol. 60, pp. 1-10).
 145. Fatema, A. H. (2016). Mycological evaluation of some ready to eat meat products with special reference to molecular characterization* Shaltout, FA;** Salem, R. M;*** El-diasty, Eman and. *Veterinary Medical Journal-Giza*, 62(3), 9-14.
 146. Fahim, A. S., Ahmed, A. A., & Eman, M. K. Heavy Metal Residues in chicken cuts up and processed chicken meat products.
 147. Shaltout, F.A.; Hanan M. Lamada, Ehsan A.M. Edris. (2020): Bacteriological examination of some ready to eat meat and chicken meals. *Biomed J Sci & Tech Res.*, 27(1): 20461- 20465.
 148. Sobhy, A., & Shaltout, F. (2020). Prevalence of some food poisoning bacteria in semi cooked chicken meat products at Kaliobia governorate with using of recent Vitek 2 compact and PCR techniques. *Benha Veterinary Medical Journal*, 38(2), 88-92.
 149. Shaltout, F. A., Riad, E. M., & AbouElhassan, A. A (2017): prevalence Of Mycobacterium Tuberculosis In Imported cattle Offals And Its lymph Nodes. *Veterinary Medical Journal-Giza (VMJG)*, 63(2), 115-122.
 150. Sobhy, A., & Shaltout, F. (2020). Detection of some food poisoning bacteria in some semi cooked chicken meat products marketed at Kaliobia governorate. *Benha Veterinary Medical Journal*, 38(2), 93-96.
 151. Shaltout, F. A. (2023). Abattoir and Bovine Tuberculosis as a reemerging foodborne disease. *Clinical Medical Reviews and Reports*, 6(1).
 152. Shaltout, F.A.(2023): Viruses in Beef, Mutton, Chevron, Venison, Fish and Poultry Meat Products. *Food Science & Nutrition Technology* 8(4):1-10.