

# More Knowledge About New Technique Used in Food Inspection

**Fahim A. Shaltout\***

Department of Veterinary Medicine, Food Control, University of Benha, Egypt.

**Corresponding Author:** Fahim A. Shaltout, Department of Veterinary Medicine, Food Control, University of Benha, Egypt.

**Received:** 📅 2024 Nov 21**Accepted:** 📅 2024 Dec 01**Published:** 📅 2024 Dec 22

## Abstract

*The Electronic noses are devices able to characterize and differentiate the aroma profiles of the various foods, especially the meat and the meat products. During the recent years advanced method the Electronic noses have been widely used in the food analysis and proved to provide a fast, simple, non expensive and non destructive method of the food assessment and quality control. The aim of this review article is to summarize the important characters of the analytic technique and to present basic fields and the typical areas of the Electronic nose use as well as the most commonly used sensor types and the patterns for the Electronic nose design. The Prospects for the future development of this technique are presented. The Methods and the researches may be a guideline for the practical Electronic nose use.*

**Keywords:** Electronic Nose, Sensors, Gas Chromatography, Odor, Food Inspection

## 1. Introduction

The most important factors affecting freshness and quality of the poultry and the fish are the color, the texture and the flavor. The Meat flavor quality is determined by composition of the VOC as its critical factor. The factors must be qualified through the electronic nose in the same way as the human senses, because these modalities are the basis for the human perception of the freshness and quality. The odor of the meat is formed by a complex mixture of the different volatile organic compounds, originating from various reactions. It is often stated that the fresh meat is almost unscented and the only the odor that can be detected by the consumer is described as bloody. Within the storage or the thermal processes, the precursors of the aroma constituents undergo the oxidation and decomposition reactions leading to a number of products which can then react further providing the organic compounds of low molecular mass and usually these secondary products are responsible for the odor development [1-7]. The characteristic aroma of the meat usually originates from the thermal processes like the cooking, the roasting or the frying. The Chemical reactions occurring during the heat treatment involve degradation of the amino acids, the peptides, the sugars, the ribonucleotides, the lipids and the vitamins. The key process in the aroma formation is Maillard reaction that occurs between reducing sugars and amino acids derivatives. The characteristic compounds for the cooked meat are the aldehydes, the ketones, the alcohols, the hydrocarbons, the pyrazine derivatives and the Sulphur compounds. A specific effect is the warmed over flavor (WOF), which develops during the storage in the meat after the cooking. It comes from the lipid oxidation induced by the iron released from the myoglobin

and hemoglobin during the cooking. The unpleasant odor is described as the metallic, the musty and the pungent and the secondary oxidation products responsible for it are hexanal, 2,3-octanediol, pentanal, 2-pentylfuran and 2-octenal. Another process connected to the odor development apart from the heat treatment of the meat is spoilage [8-14].

During the storage the bacterial activity leads to the production of the volatile organic compounds of the unpleasant odor, most of which are easily identifiable (the methanol, the ethanol, the dimethyl sulfide, the methyl thioacetate, the toluene, the nonane, the 2,3-butanediol and others). The Alcohols, the hydrocarbons, the aliphatic the ketones, the volatile acids and the benzenic compounds are the compounds typical for the physicochemical changes generated as a result of fat decomposition and the lipid oxidation [15-23]. The aroma profile of the meat can be influenced by different factors like the fat content, the animal diet, the breed, the pH, the storage condition or the heat treatment method. This subject is still not fully explored despite of many discussions. Assessment of the meat freshness can be done by analysis of mixture of volatile organic compounds. The Classical methods of analysis of the meat aroma profile involve the gas chromatography (GC) and the olfactometry (O). The fusion of these two techniques, GC-O, combines the ability of the chromatographic unit to separate different compounds and the capability of the olfactometric system to characterize them [24-32]. The chemical methods, the sensory descriptive analysis is still of great significance. The aroma profile is difficult to analyse and characterize in the laboratory since the human sense of the smell does not distinguish individual components

but rather identifies the specific mixture of volatiles as a whole. The Classical methods of the chemical analysis based on the gas chromatography result in the identification and the quantification of the particular compounds which can be considered the most important odor indicators. Often the single compounds present in meaningful quantity in the aroma profile are not noticeable by the human sense of the smell. The practice shows that there is not always a simple correlation between the concentration of particular compounds and the odor perception. The Electronic noses are analytical instruments designed to mimic the work of the human sense of the smell. In this technique the analytic process does not concentrate on the identification and quantification of the components of the mixture of volatile compounds but rather on the quantitative description of the complete aroma profile, including the relationships between its components. In many researches the time change of the profile or the profile change in relation to standard profile is important. The volatile profile is a fingerprint typical for certain test. The most important issues are presented below [33-41]. The standard Electronic noses equipment is based on a series of the gas sensors able to collect the chemical parameter coming from the headspace and to transmit it to the electronic devices.

The Different types of sensors can be used for different applications, among them the most popular are electrochemical sensors like the metal oxide semiconductors (MOS) or the conducting polymers (CP) and the piezoelectric sensors like the quartz microbalances (QCM). The Optical, calorimetric and biosensors can be also incorporated into the Electronic noses system. Those innovative types of the advanced method Electronic noses based on the gas chromatography (GC) or gas chromatography with mass spectrometry (GC/MS) have been introduced recently and their high utility has been demonstrated. the detection each peak from a mass spectrum or a chromatogram acts as a "sensor" providing also information about the chemical structure of the corresponding compound. It is a standard to couple the Electronic nose with multivariate software equipped with the software tools for the chemometric interpretation of the sensor signals [42-51]. The data collected from the sensors are analyzed using various statistical tools in order to create a numerical model of the aroma profile of the sample. When having a quantitative description of the aroma profile of the samples one can then compare the unknown samples with the reference materials or study the influence of different factors on the odor. The most effective statistical methods used in these types of investigation include principal component analysis (PCA), partial least squares regression (PLSR), The linear and canonical discriminant analysis (LDA and CDA) or the artificial neural networks (ANN). The Electronic noses are rather an unreliable tool for the determination of specific chemical compounds constituting the aroma profile, the advantages of this method over the traditional ones are conspicuous [52-61]. The analyses are fast, simple and low cost, what makes the Electronic nose a gratifying analytical method for quality control applications. The Electronic nose requires specific and time consuming training for

the staff. This review article aims to present diversity of types of the Electronic noses used for the evaluation of the meat and the meat products. The Prospects for the future development of this technique are presented. The Methods and the researches which discussed may be a guideline for the practical Electronic nose application [62-70].

## **1.1. The Applications of the Electronic Nose In the Meat Analysis**

### **1.1.1. The Spoilage Monitoring Traditional Methods of the Meat Spoilage**

The monitoring is based on measurements of chemical or biological spoilage indicators. The Standard tests include total bacterial count (TBC), rancidity measurements with thiobarbituric acid reactive substances assay (TBARS), determination of the VOCs by GC-MS and total volatile basic nitrogen (TVBN) measurements. The Colour evaluation and sensory descriptive analysis can be also useful in the detection of spoiled samples. The methods are rather costly, time consuming and require specialized staff, what makes them difficult for on line applications [71-80]. The spoilage process is strongly connected with the odor changes, the most of studies investigate the possibilities of employing the Electronic nose to this purpose. The standard procedure in this type of the study involves choosing the storage conditions (the type of the packaging and the temperature), usually mimicking storage in typical household or industrial conditions and analyzing samples after different periods of time. The importance to ensure careful preparation of the samples in order to preserve the headspace composition during storage and analysis. The sensor types used for this purpose include mainly the metal oxide semiconductors or conducting polymers. During last years a few applications using colorimetric sensors could be noticed [81-89]. The statistical tools employed for data analysis involve multivariate statistics and artificial neuron networks. The obtained results are then confronted with data from reference methods based on the microbiological evaluation, the sensory descriptive analysis or the chemical determination of the spoilage indicators. A number of fully successful attempts to employ the advanced method Electronic noses for the meat spoilage monitoring prove the utility of this analytic technique. The Electronic nose found application as a laboratory tool, yet no in industrial practice. The prospects of the method development involve further investigation of the value of information coming from particular sensors and constructing new portable instruments with reduced number of sensors in order to minimize costs and simplify the analysis [90-99].

## **1.2. The Differentiation between types of the meat**

The Determination of the animal species from which the meat was produced is an important problem of the food analysis. From the consumer's point of view, it is an essential issue not only because of the quality of the product or the meat adulteration but also because of health, dietary and religious aspects. The classical methods of the meat species identification involve expensive and time consuming molecular biology based methods or spectroscopic measurements requiring specialized equipment and

complex data analysis. The Electronic nose can be a reliable analytic tool for screening of the meat samples for the animal species origin [100-107]. The differentiation between the meat samples from sheep, cattle, poultry, and swine was performed using an electronic nose based on the gas chromatography. The simple statistical data analysis using principal component analysis (PCA) could clearly separate groups of samples derived from the different animals. The advanced method the Electronic noses, do not give information about chemical composition of the headspace, the chromatographic analysis performed by this type of the Electronic nose allowed the tentative identification of compounds responsible for the aroma profile, using Kovat's indices. The methodology presented in the study was extremely simple, fast and no expensive, providing a useful tool for the food analysis. The ability of the electronic nose to discriminate between types of the meat can be also applied in the detection of the meat adulteration with proteins from another, cheaper species. The Pork adulteration in the minced mutton was studied with an Electronic nose containing 10 metal oxide sensors. The Mutton samples were mixed with pork in different proportions and the headspace of each sample was tested with the Electronic nose. The statistic tools were implemented to analyze the data obtained (PCA, DA, PLS, MLR and BPNN) and it was clearly demonstrated that the Electronic noses methodology coupled with multivariate analysis can easily predict the degree of adulteration of pork in minced mutton. The use of the Electronic noses technique for the meat type and the quality differentiation for the meat adulteration detection and for the quality control [108-116].

### 1.3. The Evaluation of the effect of the dietary supplementation of the animals

The Animal nutrition has a potent influence on the meat quality. The Different types of dietary regimes can be applied for specific purposes, among them supplementation of functional ingredients such as vitamin E, selenium, conjugated linoleic acids or omega 3 fatty acids gained a special interest over last years. The Evaluation of the relationship between the animal diet and the meat attributes is an important feature. The investigation of the animal diet influence on overall the antioxidant power in the meat and its connection with the aroma profile. The Crossbreed steers were either live in pasture or feed on the grain and in both groups a part of the animals obtained an additional vitamin E supplementation [117,118]. The Electronic nose with 32 conducting polymer sensors was used to analyze the aroma profile of the fresh beef samples from all the four groups. The Antioxidant capacity tests were also performed. The relationship between the Electronic nose data and the antioxidant status related variables was analyzed while the linear discriminant analysis of the Electronic nose measurements was employed to investigate the meat samples grouping as a function of feeding.

The Electronic nose distinguished correctly grain and pasture produced meat as well as supplemented and non supplemented with vitamin E grain produced meat. These results show that the aroma profile of the meat is strongly

related on the antioxidant status which affects the lipids oxidation influencing the consequent production of volatile short chained aldehydes. The Electronic nose proved therefore to be a useful tool to discriminate the aroma profile of the fresh meat samples with different antioxidant potential. The influence of the animal dietary regimes on the lipid oxidation can manifest in the change of volatile compounds profile of the meat and this effect was also investigated with the use of the Electronic noses. The pigs were divided into four feeding groups with the different diets: the control diet, the supplemental vitamin E and organic selenium diet, supplemental organic selenium diet and supplemental vitamin E diet. The samples were analyzed using the electronic nose equipped with ultrafast gas chromatograph with the flame ionization detectors. The Electronic nose data were analyzed by using the AroChemBase database and the ANOVA. The measurements performed by the Electronic nose showed that the addition of the antioxidants to the pigs feed prevented the formation of the Sulphur compounds in the raw meat. The electronic nose successfully allowed to determine seventeen specific volatile compounds in the supplemented meat.

### 1.4. The Production process monitoring

There are rather few examples in the literature of direct use of the Electronic nose for production process monitoring. The Electronic nose used to identify the spoiled Iberian hams during the curing process. They discussed use of the Electronic nose for recognition of different Iberian ham ripening times. The sensors with the tin oxide semiconductor thin films were used for the tests. The sensors were doped with metal catalysts. The PCA was used for results analysis together with artificial neural network. The Electronic nose can be applied to study dynamic processes occurring during production of sausages. The investigation of the seasoning processes for the dry cured meats was performed with an electronic nose containing 12 metal oxide sensors. The Fresh pork sausages were subjected to the manufacture's protocols and 5, 7, 10, 14 days seasoning. The Research included monitoring of presence of the ochratoxin A producing and non producing *Penicillium* strains during the seasoning process. The Food contamination with the ochratoxins is very dangerous for the humans and is considered as possibly carcinogenic by The International Agency for Research on Cancer in 1993. The Electronic noses data were analyzed using the DFA. The Proposed research technique was successfully applied for the rapid prediction of the ochratoxin A. These mentioned applications of the Electronic nose are satisfactory examples of successful use of the Electronic nose based systems for quality control [119-125].

### 1.5. The Specific purposes:

The boar taint and the WOF evaluation the Boar taint and the WOF are the sensory defects in the meat flavor. The Boar taint is characteristic for pork derived from the non castrated male pigs. The boar taint is connected to the presence of the androstenone and the skatole but it was shown that the sensory evaluation of the 'boar taint' level does not always agree with the absolute concentrations of these two compounds. The need to the developing a system able to

efficiently discriminate between the different intensities of the undesirable odor. The Used Electronic nose and sensory panel to measure the intensity of the boar taint in the entire male pigs. The electronic nose based on ion mobility spectrometry was employed to mimic the responses given by the sensory panel. The data from the Electronic noses analysis were calibrated using canonical correlation with the sensory measurement and a discriminant function for separating levels of the boar taint in the pork by the Electronic nose was developed [126-132]. The Sensorics research affirmed stronger correlation of the boar taint with the androsterone than with the skatole. The research showed that the Electronic nose technique based on the ion mobility spectrometry may have a potential for a rapid sorting of boar fat at the slaughter line. The warmed over flavor develops in the meat which has been pre cooked, chill stored and reheated. The sensory analysis of volatile compounds on the meatballs derived from the pigs fed with standard diet supplemented with the addition of rapeseed and palm oil using solid state based gas sensor array system (the Electronic nose) and the gas chromatography/gas spectrometry together with measurements of the thiobarbituric acid reactive substances (TBARS). The Obtained data were analyzed using partial least square regression modelling (PLSR). The MOS sensor responses showed to be significantly related to WOF characteristics detected by both the sensory and the chemical analysis. This shows the potential of using gas sensor technology to monitor WOF in the pork. The simple and rapid method of prediction of WOF in the cooked chicken by colorimetric sensor array. The Data from colorimetric sensor array was classified using principal component analysis and hierarchical cluster analysis. The Research showed that colorimetric sensor array may be successfully used to predict WOF development in the cooked chicken meat [133-138].

## 2. Conclusions

The electronic noses provide a fast, simple and non destructive method of the meat analysis. The Electronic noses were successfully employed mainly in the quality control of the meat being able to monitor spoilage or adulterations and the obtained results were in accordance with the sensory evaluation, offering a reliable tool for on line analysis. The ability of the Electronic nose to determine the antioxidant status of the meat samples was also proved. It can be stated that every factor that influences the aroma profile of the meat, could be potentially indirectly analyzed with the Electronic nose system and this field still seems to be insufficiently explored. The development of new types of advanced method Electronic noses, based on gas chromatography, opens new perspectives for analysis of the aroma profile of the meat. Despite of many scientific works proving usefulness of the Electronic nose based on different types of sensors or GC techniques for the meat quality appraisal or spoilage detection, use of the Electronic nose in industry practice is insufficient. There was not found any description or report on large scale industrial application of Electronic nose. Almost every paper describes potentially large possibilities of use of Electronic nose in industrial

practice, but none reports real life implementation. This may be explained by sensor vulnerability (sensor time drift), relatively the high Electronic nose costs and effort consuming staff training. The above does not mean that the Electronic nose will not be used widely in future in industrial practice. There is still lack of effective application, although the potential of the Electronic nose approach was proven.

## References

1. AbouElhassan, A. (2017). Prevalence of Mycobacterium Tuberculosis in Imported cattle Offals and its lymph Nodes. *Veterinary Medical Journal (Giza)*, 63(2), 115-122.
2. El Barbri, N., Llobet, E., El Bari, N., Correig, X., & Bouchikhi, B. (2008). Electronic nose based on metal oxide semiconductor sensors as an alternative technique for the spoilage classification of red meat. *Sensors*, 8(1), 142-156.
3. Shaltout, F. A., Riad, E. M., & Asmaa, A. E. PREVALENCE OF MYCOBACTERIUM SPP. IN CATTLE MEAT AND OFFALS SLAUGHTERED IN AND OUT ABATTOIR.
4. Abd Elaziz, O., Hassanin, F., Shaltout, F., & Mohammed, O. (2021). Prevalence of some foodborne parasitic affection in slaughtered animals in local Egyptian abattoir. *Benha Veterinary Medical Journal*, 41(1), 111-114.
5. Abd El-Aziz, O. M., Hassanin, F. S., Shaltout, F. A. E., & Mohammed, O. (2021). Prevalence of some zoonotic parasitic affections in sheep carcasses in a local abattoir in Cairo, Egypt. *Benha Veterinary Medical Journal*, 41(1), 115-119.
6. Jónsdóttir, R., Ólafsdóttir, G., Chanie, E., & Haugen, J. E. (2008). Volatile compounds suitable for rapid detection as quality indicators of cold smoked salmon (*Salmo salar*). *Food chemistry*, 109(1), 184-195.
7. Al Shorman, A. A. M., Shaltout, F., and Hilat, N., (1999). Detection of certain hormone residues in meat marketed in Jordan. *Jordan University of Science and Technology, 1st International Conference on Sheep and goat Diseases and Productivity*, 23-25 October, 1999.
8. Saleh, E., Shaltout, F., & Abd Elaal, E. (2021). Effect of some organic acids on microbial quality of dressed cattle carcasses in Damietta abattoirs, Egypt. *Damanhour Journal of Veterinary Sciences*, 5(2), 17-20.
9. Khan, M. I., Jo, C., & Tariq, M. R. (2015). Meat flavor precursors and factors influencing flavor precursors—A systematic review. *Meat science*, 110, 278-284.
10. Shaltout, F. A. E. (2017). Microbiological evaluation of some heat treated fish products in Egyptian markets. *EC Nutrition*, 12, 124-132.
11. Edris, A., Hassan, M. A., Shaltout, F. A., & Elhosseiny, S. (2013). Chemical evaluation of cattle and camel meat. *Benha veterinary medical journal*, 24(2), 191-197.
12. Edris, A. M., Hassan, M. A., Shaltout, F. A., & Elhosseiny, S. (2012). Detection of *E. coli* and *Salmonella* organisms in cattle and camel meat. *Benha veterinary medical journal*, 24(2), 198-204.
13. Edris, A. M., Hemmat, M. I., Shaltout, F. A., Elshater, M. A., & Eman, F. M. I. (2012). Study on incipient spoilage of chilled chicken cuts-up. *Benha veterinary medical*

- journal, 23(1), 81-86.
14. Edris, A. M., Hemmat, M. I., Shaltout, F. A., Elshater, M. A., & Eman, F. M. I. (2012). Chemical analysis of chicken meat with relation to its quality. *Benha veterinary medical Journal*, 23(1), 87-92.
  15. Edris, A.M.; Shaltout, F. and Abd Allah, A.M. (2005): Incidence of *Bacillus cereus* in some meat products and the effect of cooking on its survival. *Zag. Vet. J* 33 (2):118-124.
  16. Edris, A. M., Shaltout, F. A., & Arab, W. S. (2005). Bacterial evaluation of quail Meat. *Benha Vet. Med. J*, 16(1), 1-14.
  17. Edris, A. M., Shaltout, F., Salem, G. H., & El-Toukhy, E. I. (2011). Incidence and isolation of *Salmonellae* from some meat products. Benha University, Faculty of Veterinary Medicine. In Fourth Scientific Conference (pp. 172-179).
  18. Shaltout, F. A. E. (2017). Microbiological evaluation of some heat treated fish products in Egyptian markets. *EC Nutrition*, 12, 124-132.
  19. Leroy, F., Vasilopoulos, C., Van Hemelryck, S., Falony, G., & De Vuyst, L. (2009). Volatile analysis of spoiled, artisan-type, modified-atmosphere-packaged cooked ham stored under different temperatures. *Food Microbiology*, 26(1), 94-102.
  20. Edris, A. M., Shaltout, F. A., Salem, G. H., & El-Toukhy, E. I. (2011). Plasmid profile analysis of *Salmonellae* isolated from some meat products. Benha University, Faculty of Veterinary Medicine. In Fourth Scientific Conference (pp. 194-201).
  21. Ragab, A., Edris, A. M., Shaltout, F. A., & Salem, A. M. (2022). Effect of titanium dioxide nanoparticles and thyme essential oil on the quality of the chicken fillet. *Benha veterinary medical journal*, 41(2), 38-40.
  22. Hassan, M. A., Shaltout, F. A., Arfa, M. M., Mansour, A. H., & Saudi, K. R. (2013). Biochemical studies on rabbit meat related to some diseases. *Benha Vet. Med. J*, 25(1), 88-93.
  23. Hassan, M.A., Shaltout, F., (1997). Occurrence of Some Food Poisoning Microorganisms In Rabbit Carcasses. *Alex.J.Vet.Science*, 13(1), 55-61.
  24. Hassan, M., Shaltout, F. A., & Saqur, N. (2020). Histamine in some fish products. *Archives of Animal Husbandry & Dairy Science*, 2(1), 1-3.
  25. Hassan, M. A., Shaltout, F. A., (2004). Comparative Study on Storage Stability of Beef, Chicken meat, and Fish at Chilling Temperature. *Alex. J. Vet. Science*, 20(21), 21-30.
  26. Hassan, M. A., Shaltout, F. A., Arfa, M. M., Mansour, A. H., & Saudi, K. R. (2013). Biochemical studies on rabbit meat related to some diseases. *Benha Vet. Med. J*, 25(1), 88-93.
  27. Rivas-Cañedo, A., Juez-Ojeda, C., Nuñez, M., & Fernández-García, E. (2011). Effects of high-pressure processing on the volatile compounds of sliced cooked pork shoulder during refrigerated storage. *Food chemistry*, 124(3), 749-758.
  28. Hassan, M. A., Shaltout, F. A., Maarouf, A. A., & El-Shafey, W. S. (2014). Psychrotrophic bacteria in frozen fish with special reference to *Pseudomonas* species. *Benha Vet. Med. J*, 27(1), 78-83.
  29. Hassan, M. A., Shaltout, F. A., Arfa, M. M., Mansour, A. H., & Saudi, K. R. (2013). Biochemical studies on rabbit meat related to some diseases. *Benha Vet. Med. J*, 25(1), 88-93.
  30. Hassanin, F. S., Hassan, M. A., Shaltout, F. A., Shawqy, N. A., & Abd-Elhameed, G. A. (2017). Chemical criteria of chicken meat. *Benha veterinary medical journal*, 33(2), 457-464.
  31. Hassanien-faten, S., Hassan, M. A., Shaltout, S., & Elrais-Amina, M. *CLOSTRIDIUM PERFRINGENS* IN VACUUM PACKAGED MEAT PRODUCTS.
  32. Hassanien, F., Shaltout, F. A. E., Fahmey, M. Z., & Elsukkary, H. F. A. (2020). Bacteriological quality guides in local and imported beef and their relation to public health. *Benha Veterinary Medical Journal*, 39(1), 125-129.
  33. Hassanin, F. S., Shaltout, F. A., & Mostafa, E. M. (2013). Parasitic affections in edible offal. *Benha Vet. Med. J*, 25(2), 34-39.
  34. Hassanin, F. S., Shaltout, F. A., Lamada, H. M., & Abd Allah, E. M. (2011). The effect of preservative (nisin) on the survival of *Listeria monocytogenes*. *Benha veterinary medical journal (2011)-special issue*, 141-145.
  35. Khattab, E., Shaltout, F., & Sabik, I. *Benha Veterinary Medical Journal*.
  36. Saad, S. M., Shaltout, F. A., Farag, A. A., & Mohammed, H. F. (2022). Organophosphorus Residues in Fish in Rural Areas. *Journal of Progress in Engineering and Physical Science*, 1(1), 27-31.
  37. Saif, M., Saad, S., Shaltout, F., Hassanin, F. S., & Zaghloul, M. (2019). Molecular detection of enterotoxigenic *Staphylococcus aureus* in ready to eat beef products. *Benha Veterinary Medical Journal*, 37(1), 7-11.
  38. Saif, M., Saad, S., Hassanin, F. S., Shaltout, F., & Zaghloul, M. (2019). Prevalence of methicillin-resistant *Staphylococcus aureus* in some ready-to-eat meat products. *Benha Veterinary Medical Journal*, 37(1), 12-15.
  39. Dissing, B. S., Papadopoulou, O. S., Tassou, C., Ersbøll, B. K., Carstensen, J. M., Panagou, E. Z., & Nychas, G. J. (2013). Using multispectral imaging for spoilage detection of pork meat. *Food and Bioprocess Technology*, 6(9), 2268-2279.
  40. Abdelmordy, A., Shaltout, F., & Saad, S. M. (2023). Studies on Pesticides Residues in Fish in Menofia Governorate. *Benha Journal of Applied Sciences*, 8(5), 323-330.
  41. Abdelmordy, A., Shaltout, F., & Saad, S. M. (2023). Organochlorine Residues in Fish in Rural Areas. *Benha Journal of Applied Sciences*, 8(5), 331-336.
  42. Shaltout, F., Hussein, M., & khaled Elsayed, N. (2023). Histological Detection of Unauthorized Herbal and Animal Contents in Some Meat Products. *Journal of Advanced Veterinary Research*, 13(2), 157-160.
  43. Ghanem, A., Shaltout, F., & Heikal, G. I. (2022). Mycological quality of some chicken meat cuts in Gharbiya governorate with special reference to *Aspergillus flavus* virulent factors. *Benha Veterinary medical journal*, 42(1), 12-16.
  44. Darwish, W., Shaltout, F. A., Salem, R. M., Eldiasty, E. M., & Diab, F. A. (2022). Seasonal Impact on the Prevalence of Yeast Contamination of Chicken Meat Products and Edible Giblets. *Journal of Advanced Veterinary Research*, 12(5), 641-644.
  45. Shaltout, F. A., Barr, A. A. H., & Abdelaziz, M. E. (2022). Pathogenic microorganisms in meat products.

- Biomedical Journal of Scientific & Technical Research, 41(4), 32836-32843.
46. Shaltout, F. A., & Koura, H. A. (2017). Impact of some essential oils on the quality aspect and shelf life of meat. *Benha veterinary medical journal*, 33(2), 351-364.
  47. Mohammed, I. Z. (2020). Bacteriological profile of some raw chicken meat cuts in Ismailia city, Egypt. *Benha Veterinary Medical Journal*, 39(1), 11-15.
  48. Afify, S. A., Shaltout, F., & Mohammed, I. Z. (2020). Detection of *E. coli* O157 and *Salmonella* species in some raw chicken meat cuts in Ismailia province, Egypt. *Benha Veterinary Medical Journal*, 39(1), 101-104.
  49. Shaltout, F. A., El-diasty, E. M., & Asmaa-Hassan, M. A. (2019). Hygienic quality of ready to eat cooked meat in restaurants at cairo governorate. *Journal of Global Biosciences*, 8(12), 6627-6641.
  50. Hong, X., Wang, J., & Hai, Z. (2012). Discrimination and prediction of multiple beef freshness indexes based on electronic nose. *Sensors and Actuators B: Chemical*, 161(1), 381-389.
  51. Shaltout, F. A., Nasief, M. Z., Lotfy, L. M., & Gamil, B. T. *Benha Veterinary Medical Journal*.
  52. Shaltout, F. (2019). Poultry Meat. *Scholarly Journal of Food and Nutrition*, 22 1-2.
  53. Shaltout, F. (2019). Food Hygiene and Control. *Food Science and Nutrition Technology* 4(5), 1-2.
  54. Arakeeb, S. M., Hassanien, F., Shaltout, F., & Homouda, S. (2019). Natural preservatives in raw chicken meat. *Benha Veterinary Medical Journal*, 37(1), 41-45.
  55. del Olmo, A., Calzada, J., & Nuñez, M. (2014). Effect of high-pressure-processing and modified-atmosphere-packaging on the volatile compounds and odour characteristics of sliced ready-to-eat "lacón", a cured-cooked pork meat product. *Innovative Food Science & Emerging Technologies*, 26, 134-142.
  56. Hazaa, W., Shaltout, F., & El-Shater, M. A. (2019). Prevalence of some chemical hazards in some meat products. *Benha Veterinary Medical Journal*, 37(1), 32-36.
  57. Hazaa, W., Shaltout, F., & El-Shater, M. (2019). Identification of Some Biological Hazards in Some Meat Products. *Benha Veterinary Medical Journal*, 37(1), 27-31.
  58. Gaafar, R., Hassanin, F. S., Shaltout, F., & Zaghloul, M. (2019). Molecular detection of enterotoxigenic *Staphylococcus aureus* in some ready to eat meat based sandwiches. *Benha Veterinary Medical Journal*, 37(1), 22-26.
  59. Gaafar, R., Hassanin, F. S., Shaltout, F., & Zaghloul, M. (2019). Hygienic profile of some ready to eat meat product sandwiches sold in Benha city, Qalubiya Governorate, Egypt. *Benha Veterinary Medical Journal*, 37(1), 16-21.
  60. Saad, S. M., Shaltout, F. A., Abou Elroos, N. A., & El-nahas, S. B. (2019). Antimicrobial effect of some essential oils on some pathogenic bacteria in minced meat. *J Food Sci Nutr Res*, 2(1), 013-021.
  61. Kim, S. Y., Li, J., Lim, N. R., Kang, B. S., & Park, H. J. (2016). Prediction of warmed-over flavour development in cooked chicken by colorimetric sensor array. *Food Chemistry*, 211, 440-447.
  62. Shaltout, F. A., Saad, M., Abo El-Roos, N., & Saber, E. (2019). Incidence of *Staphylococci* and *E. coli* in Meat and Some Meat Products. *EC Nutrition*, 14(6), 171-179.
  63. Saad, S. M., Hassanin, F. S., Shaltout, F. A., Marionette, Z. N., & Marwa, Z. S. (2019). Prevalence of methicillin-resistant *Staphylococcus aureus* in some ready-to-eat meat products. *American Journal of Biomedical Science & Research*, 4(6), 460-464.
  64. Shaltout, F. (2019). Pollution of Chicken Meat and Its Products by Heavy Metals. *Research and Reviews on Healthcare: Open Access Journal*, 4(3), 381-3382.
  65. Chen, Q., Hui, Z., Zhao, J., & Ouyang, Q. (2014). Evaluation of chicken freshness using a low-cost colorimetric sensor array with AdaBoost-OLDA classification algorithm. *LWT-Food Science and Technology*, 57(2), 502-507.
  66. Shaltout, F. A., EL-diasty, E. M., & Mohamed, M. S. (2018). Effects of chitosan on quality attributes fresh meat slices stored at 40C. *Benha veterinary medical journal*, 35(2), 157-168.
  67. Shaltout, F., & Abdel-Aziz, A. (2004). *Salmonella enterica* serovar Enteritidis in poultry meat and their epidemiology. *Veterinary Medical Journal (Giza)*, 52(3), 429-436.
  68. Shaltout, F. A., El-Shorah, H. F., El Zahaby, D. I., & Lotfy, L. M. (2018). Bacteriological profile of chicken meat products. *Food Nutr Current Res*, 1(3), 83-90.
  69. Shaltout, F., El-Shater, M. A., & Abd El-Aziz, W. M. (2015). Bacteriological assessment of street vended meat products sandwiches in Kalyobia Governorate. *Benha veterinary medical journal*, 28(2), 58-66.
  70. Fahim, H. M. (2019). Evaluate antibiotic residues in beef and effect of cooking and freezing on it. *Benha Veterinary Medical Journal*, 36(2), 109-116.
  71. Shaltout, F. A., Zakaria, I. M., & Nabil, M. E. (2018). Incidence of some anaerobic bacteria isolated from chicken meat products with special reference to *Clostridium perfringens*. *Nutrition and Food Toxicology*, 2(5), 429-438.
  72. Shaltout, F. A., Maarouf, A. A., & Elkhoully, M. E. (2017). Bacteriological Evaluation of Frozen Sausage. *Nutrition and Food Toxicology*, 1(5), 174-185.
  73. Shaltout, F. A., El-Toukhy, E. I., & Abd El-Hai, M. M. (2019). Molecular Diagnosis of *Salmonellae* in Frozen Meat and Some Meat Products. *Nutrition and Food Technology Open Access*, 5(1), 1-6.
  74. Shaltout, F. A., Ali, A. M., & Rashad, S. M. (2016). Bacterial Contamination of Fast Foods. *Benha Journal of Applied Sciences (BJAS)*, 1(2), 45-51.
  75. Shaltout, F. A., Eltanani, J., & Elmelegy, A. S. (2015). Microbiological status of meat and chicken received to University student hostel. *Benha veterinary medical journal*, 29(2), 187-192.
  76. Casaburi, A., Piombino, P., Nychas, G. J., Villani, F., & Ercolini, D. (2015). Bacterial populations and the volatilome associated to meat spoilage. *Food microbiology*, 45, 83-102.
  77. Saad, S. M., Edris, A. M., Shaltout, F. A., & Edris, S. (2012). Isolation and identification of *salmonellae* and *E. coli* from meat and poultry cuts by using A. multiplex PCR.

- Benha Vet. Med. J. special, (16-26).
78. Saad, S. M., & Shaltout, F. A. (1998). Mycological Evaluation of camel carcasses at Kalyobia Abattoirs. *Veterinary Medical Journal (Giza)*, 46(3), 223-229.
  79. Saad, S. M., Shaltout, F. A., Abou Elroos, N. A., & El-nahas, S. B. (2019). Antimicrobial effect of some essential oils on some pathogenic bacteria in minced meat. *J Food Sci Nutr Res*, 2(1), 013-021.
  80. Saad, S. M., Hassanin, F. S., Shaltout, F. A., Marionette, Z. N., & Marwa, Z. S. (2019). Prevalence of methicillin-resistant *Staphylococcus aureus* in some ready-to-eat meat products. *American Journal of Biomedical Science & Research*, 4(6), 460-464.
  81. Shaltout, F. A., Saad, M., Abo El-Roos, N., & Saber, E. (2019). Incidence of *Staphylococci* and *E. coli* in Meat and Some Meat Products. *EC Nutrition*, 14(6), 171-179.
  82. Shaltout, F. A., Riad, E. M., TES, A., & AbouElhassan, A. (2017). Studying the Effect of Gamma Irradiation on Bovine Offal's Infected with *Mycobacterium tuberculosis* Bovine Type. *Journal of Food Biotechnology Research*, 1(6), 1-5.
  83. Shaltout, F. A., Maarouf, A. A., & Elkhoully, M. E. (2017). Bacteriological Evaluation of Frozen Sausage. *Nutrition and Food Toxicology*, 1(5), 174-185.
  84. Shaltout, F. A., Zakaria, I. M., & Nabil, M. E. (2018). Incidence of some anaerobic bacteria isolated from chicken meat products with special reference to *Clostridium perfringens*. *Nutrition and Food Toxicology*, 2(5), 429-438.
  85. Biniecka, M., & Caroli, S. (2011). Analytical methods for the quantification of volatile aromatic compounds. *TrAC Trends in Analytical Chemistry*, 30(11), 1756-1770.
  86. Faheim, A. S., Mohamed, A. H., & Fatin, S. H. (2004). Thermal inactivation of enterohaemorrhagic *Escherichia coli* O157:h7 and its sensitivity to nisin and lactic acid cultures. 1st Ann Confr FVM Moshtohor.
  87. Aziz, S., Mahmoud, E. D., El-mesalamy, M., & El-shaer, M. (2014). Study on fungal contamination of some chicken meat products with special reference to the use of PCR for its identification. *Veterinary Medical Journal (Giza)*, 60(2), 1-22.
  88. Shaltout, F. A. (2002). Microbiological aspects of semi-cooked chicken meat products. *Benha Vet. Med. J.*, 13(2), 15-26.
  89. Shaltout, F. A., & Koura, H. A. (2017). Impact of some essential oils on the quality aspect and shelf life of meat. *Benha veterinary medical journal*, 33(2), 351-364.
  90. Shaltout, F. A. E., Farouk, M., Ibrahim, H. A., & Afifi, M. E. (2017). Incidence of Coliform and *Staphylococcus aureus* in ready to eat fast foods. *Benha veterinary medical journal*, 32(1), 13-17.
  91. Shaltout, F. A., Zakaria, I. M., & Nabil, M. E. (2017). Detection and typing of *Clostridium perfringens* in some retail chicken meat products. *Benha veterinary medical journal*, 33(2), 283-291.
  92. Shaltout, F. A. M. (1992). Studies on mycotoxins in meat and meat by-products.
  93. Shaltout, F. A. M. (1996). Mycological and Mycotoxicological profile of Some Meat products.
  94. Huang, X. W., Zou, X. B., Shi, J. Y., Guo, Y., Zhao, J. W., Zhang, J., & Hao, L. (2014). Determination of pork spoilage by colorimetric gas sensor array based on natural pigments. *Food Chemistry*, 145, 549-554.
  95. Shaltout, F. A. (1998). Proteolytic Psychrotrophes in Some Meat products. *Alex. Vet. Med. J.*, 14(2), 97-107.
  96. Shaltout, F. A. (1999). Anaerobic Bacteria in Vacuum Packed Meat Products. *Benha Vet. Med. J.*, 10(1), 1-10.
  97. Shaltout, F. A. (2000). PROTOZOAL F OODBORNE PATHOGENS IN SOME MEAT PRODCUTS. *Assiut Veterinary Medical Journal*, 42(84), 54-59.
  98. Shaltout, F. A. (2001). Quality evaluation of sheep carcasses slaughtered at Kalyobia abattoirs. *Assiut Veterinary Medical Journal*, 46(91), 150-159.
  99. Shaltout, F. A. (2002). Microbiological aspects of semi-cooked chicken meat products. *Benha Vet. Med. J.*, 13(2), 15-26.
  100. Calkins, C. R., & Hodgen, J. M. (2007). A fresh look at meat flavor. *Meat science*, 77(1), 63-80.
  101. Shaltout, F. A. (2003, April). *Yersinia Enterocolitica* in some meat products and fish marketed at Benha city. In *The Third international conference Mansoura (Vol. 1, pp. 29-30)*.
  102. Fahim, A. S. (2020). Microbiological Quality of Chicken Carcasses at Modern Poultry Plant. *J. Nutrition and Food Processing*, 3(1).
  103. Shaltout, F., & Abdel-Aziz, A. (2004). *Salmonella enterica* serovar *Enteritidis* in poultry meat and their epidemiology. *Veterinary Medical Journal (Giza)*, 52(3), 429-436.
  104. Shaltout, F. A., & Abdel Aziz, A. M. (2004). *Escherichia coli* strains in slaughtered animals and their public health importance. *J. Egypt. Vet. Med. Association*, 64(2), 7-21.
  105. Shaltout, F. A., Amin, R. A., Nassif, M. Z., & Abdelwahab, S. A. (2014). Detection of aflatoxins in some meat products. *Benha Vet Med J*, 27(2), 368-374.
  106. Shaltout, F. A., & Afify, J. R. EM and Abo Elhasan, Asmaa, A. (2012): Improvement of microbiological status of oriental sausage. *Journal of Egyptian Veterinary Medical Association*, 72(2), 157-167.
  107. Shaltout, F. A. (2023). Chemical analytical studies on rabbit meat and liver. *Benha Veterinary Medical Journal (Egypt)*, 7(2).
  108. Shaltout, F. A., & Edris, A. M. (1999). Contamination of shawerma with pathogenic yeasts. *Assiut Veterinary Medical Journal*, 41(81), 170-176.
  109. Shaltout, F. A., Eldiasty, E. M., & Mohamed, M. S. (2014, June). Incidence of lipolytic and proteolytic fungi in some chicken meat products and their public health significance. In *1st Scientific conference of food safety and Technology (pp. 79-89)*.
  110. Hassan, A. (2016). Mycological quality of chicken carcasses and extending shelf-life by using preservatives at refrigerated storage. *Veterinary Medical Journal (Giza)*, 62(3), 1-7.
  111. Diab, F. (2016). Mycological evaluation of some ready to eat meat products with special reference to molecular characterization. *Veterinary Medical Journal*

- (Giza), 62(3), 9-14.
112. Shaltout, F., El-Shater, M. A., & Abd El-Aziz, W. M. (2015). Bacteriological assessment of street vended meat products sandwiches in Kalyobia Governorate. *Benha veterinary medical journal*, 28(2), 58-66.
113. Shewail, A., SHALTOUT, F., & GERGES, T. (2018). Impact of some organic acids and their salts on microbial quality and shelf life of beef. *Assiut veterinary medical journal*, 64(159), 164-177.
114. Shaltout, F. A., Ghoneim, A. M., Essmail, M. E., & Yousseif, A. (2001). Studies on aflatoxin B1 residues in rabbits and their pathological effects. *J. Egypt. Vet. Med. Association*, 61(2), 85-103.
115. Acevedo, C. A., Creixell, W., Pavez-Barra, C., Sánchez, E., Albornoz, F., & Young, M. E. (2012). Modeling volatile organic compounds released by bovine fresh meat using an integration of solid phase microextraction and databases. *Food and Bioprocess Technology*, 5(6), 2557-2567.
116. Shaltout, F. A., & El-Lawendy, H. M. (2003). Heavy Metal Residues in Shawerma. *Beni-Suef Vet. Med J*, 13, 213-224.
117. Shaltout, F. A., & Hashim, M. F. (2002). Histamine in salted, Smoked and Canned Fish products. *Benha Vet. Med. J*, 13(1), 1-11.
118. Fahim, S. A., Mohammed, H. F., & Saber, E. N. (2015). Detection of some heavy metals in fish (*Tilapia nilotica* and *Clarias lazera*) at Menofia governorate. *Benha Veterinary Medical Journal*, 29(1), 124-128.
119. Shaltout, F. A., & Ibrahim, H. M. (1997). Quality evaluation of luncheon and Alexandrian sausage. *Benha Vet. Med. J*, 10(1), 1-10.
120. Shaltout, F. A., Marionette, Z. N., & Shakran, A. M. (2014). Quality of battered and breaded chicken meat products.
121. Huang, L., Zhao, J., Chen, Q., & Zhang, Y. (2014). Nondestructive measurement of total volatile basic nitrogen (TVB-N) in pork meat by integrating near infrared spectroscopy, computer vision and electronic nose techniques. *Food chemistry*, 145, 228-236.
122. Shaltout, F. A., Amani, M. S., Mahmoud, A. H., & Abd Elraheem, K. A. (2013). Bacterial aspect of cooked meat and edible offal at street vendors level. *J. Benha vet. Med*, 24(1), 320-328.
123. Shaltout, F. A., & Salem, R. M. T. (2000). Moulds, aflatoxin B1 and ochratoxin A in forzen livers and meat products.
124. Al-Tarazi, Y., Al-Zamil, A., Shaltout, F., & Abdel-Samei, H. (2003). Sanitary status of raw cow milk marketed in northern Jordan. *Assiut Veterinary Medical Journal*, 49(96), 180-194.
125. Shaltout, F. A., Zakaria, I. M., & Nabil, M. E. (2018). Incidence of some anaerobic bacteria isolated from chicken meat products with special reference to *Clostridium perfringens*. *Nutrition and Food Toxicology*, 2(5), 429-438.
126. Shaltout, F. A., Eldiasty, E. M., & Mohamed, M. S. (2014, June). Incidence of lipolytic and proteolytic fungi in some chicken meat products and their public health significance. In 1st Scientific conference of food safety and Technology (pp. 79-89).
127. Hassan, A. (2016). Mycological quality of chicken carcasses and extending shelf-life by using preservatives at refrigerated storage. *Veterinary Medical Journal (Giza)*, 62(3), 1-7.
128. Shaltout, F. A., Salem, R. M., El-Diasty, E. M., & Hassan, W. I. M. (2019). Effect of Lemon fruits and turmeric extracts on fungal pathogens in refrigerated chicken fillet meat. *Global Veterinaria*, 21(3), 156-160.
129. Aziz, S., Mahmoud, E. D., El-mesalamy, M., & El-shaer, M. (2014). Study on fungal contamination of some chicken meat products with special reference to the use of PCR for its identification. *Veterinary Medical Journal (Giza)*, 60(2), 1-22.
130. Diab, F. (2016). Mycological evaluation of some ready to eat meat products with special reference to molecular characterization. *Veterinary Medical Journal (Giza)*, 62(3), 9-14.
131. Fahim, A. S., Ahmed, A. A., & Eman, M. K. Heavy Metal Residues in chicken cuts up and processed chicken meat products.
132. Jaffrès, E., Lalanne, V., Mace, S., Cornet, J., Cardinal, M., Sérot, T., ... & Joffraud, J. J. (2011). Sensory characteristics of spoilage and volatile compounds associated with bacteria isolated from cooked and peeled tropical shrimps using SPME-GC-MS analysis. *International Journal of Food Microbiology*, 147(3), 195-202.
133. Shaltout, F. A., Lamada, H. M., & Edris, E. A. (2020). Bacteriological examination of some ready to eat meat and chicken meals.
134. Sobhy, A., & Shaltout, F. *Benha Veterinary Medical Journal*.
135. Sobhy, A., & Shaltout, F. (2020). Detection of some food poisoning bacteria in some semi cooked chicken meat products marketed at Kaliobyia governorate. *Benha Veterinary Medical Journal*, 38(2), 93-96.
136. Descalzo, A. M., Rossetti, L., Grigioni, G., Irurueta, M., Sancho, A. M., Carrete, J., & Pensel, N. A. (2007). Antioxidant status and odour profile in fresh beef from pasture or grain-fed cattle. *Meat Science*, 75(2), 299-307.
137. Shaltout, F. A. (2023). Abattoir and Bovine Tuberculosis as a reemerging foodborne disease. *Clinical Medical Reviews and Reports*, 6(1).
138. Shaltout, F. A. (2023). Viruses in Beef, Mutton, Chevon, Venison, Fish and Poultry Meat Products. *Mathews Journal of Veterinary Science*, 7(5), 1-12.